

Fallopian Tube (C57.0)

A. Summary Stage Rule Changes

1. Involvement of:
 - inguinal lymph nodes
 - lateral sacral (laterosacral) lymph nodes
 - Distant (SS1977) to Regional lymph nodes (SS2000)*
2. Extension to:
 - cul de sac (rectouterine pouch)
 - omentum
 - ovary, contralateral
 - rectosigmoid
 - sigmoid
 - small intestine.
 - Distant (SS1977) to Regional by direct extension only (SS2000)*

B. Stage Shift

<u>Stage</u>	<u>SS 1977</u>	<u>SS 2000</u>
Localized	33.9%	33.9%
Regional-Ext	6.7%	17.7%
Regional-Node	4.3%	4.9%
Reg-Ext + Node	1.5%	3.7%
Distant	51.7%	37.9%
Unknown	1.8%	1.8%
Total	100.0%	100.0%

	} 12.5%		
		} 26.3%	

C. Data Interpretation

Summary Stage 2000 is not compatible with Summary Stage 1977 codes. The major shift is cases in distant stage according to SS1977 moving to a regional stage in SS2000.

Indirect Coding to Summary Stage 1977 and Summary Stage 2000 based on Extent of Disease Codes
SEER Program Data, 1998-2001, Fallopian Tube

ss1977

ss2000

	0: In Situ	1: Localized	2: Regional	3: Reg LN	4: Reg + LN	7: Distant	9: Unknown	Total
1: Localized		111	0	0	0	0	0	111
2: Regional		0	22	0	0	0	0	22
3: Reg LN		0	0	14	0	0	0	14
4: Reg + LN		0	0	0	5	0	0	5
7: Distant		0	36	2	7	124	0	169
9: Unknown		0	0	0	0	0	6	6
Total		111	58	16	12	124	6	327

Proportion:	0: In Situ	1: Localized	2: Regional	3: Reg LN	4: Reg + LN	7: Distant	9: Unknown	Total
1: Localized		0.33945	0	0	0	0	0	0.33945
2: Regional		0	0.067278	0	0	0	0	0.067278
3: Reg LN		0	0	0.042813	0	0	0	0.042813
4: Reg + LN		0	0	0	0.015291	0	0	0.015291
7: Distant		0	0.110092	0.006116	0.021407	0.379205	0	0.51682
9: Unknown		0	0	0	0	0	0.018349	0.018349
Total		0.33945	0.17737	0.04893	0.036697	0.379205	0.018349	1

p0 0.862385
pe 0.326132
K 0.795784

