

Average Annual Age-adjusted (2000 U.S. Standard) NAACCR (U.S.) Combined² Cancer Incidence Rates¹ and Rate Ratios for Selected Cancer Sites³ by Race/Ethnicity and Gender, 2010-2014

Males

Cancer Site	Rates			Rate Ratios		
	Hispanic	Non-Hispanic White	Non-Hispanic Black	Hispanic:Non-Hispanic White	Hispanic:Non-Hispanic Black	Hispanic:Non-Hispanic Black
All Sites	386.3	510.8	561.0	0.76		0.69
Prostate	97.1	107.1	186.8	0.91		0.52
Colon and Rectum	41.9	45.2	56.4	0.93		0.74
Lung and Bronchus	40.6	75.9	88.0	0.53		0.46
Kidney and Renal Pelvis	20.8	22.1	24.8	0.94		0.84
NH ⁴ Lymphoma	19.8	23.9	17.2	0.83		1.15
Liver and IBD ⁴	19.7	10.0	17.2	1.97		1.15
Urinary Bladder	19.5	39.8	19.8	0.49		0.98
Leukemia	13.1	18.4	13.7	0.71		0.96
Stomach	12.9	7.8	14.3	1.65		0.90
Pancreas	12.0	14.3	17.3	0.84		0.69
Oral Cavity and Pharynx	10.5	18.8	14.8	0.56		0.71

Females

Cancer Site	Rates			Rate Ratios		
	Hispanic	Non-Hispanic White	Non-Hispanic Black	Hispanic:Non-Hispanic White	Hispanic:Non-Hispanic Black	Hispanic:Non-Hispanic Black
All Sites	329.6	436.0	407.5	0.76		0.81
Breast	91.9	128.7	125.5	0.71		0.73
Colon and Rectum	29.3	34.5	41.7	0.85		0.70
Lung and Bronchus	25.2	57.5	50.1	0.44		0.50
Corpus and Uterus, NOS ⁴	22.2	26.6	25.6	0.83		0.87
Thyroid	20.2	22.7	13.8	0.89		1.46
NH ⁴ Lymphoma	15.1	16.4	12.1	0.92		1.25
Kidney and Renal Pelvis	12.0	11.3	12.9	1.06		0.93
Ovary	10.3	12.0	9.4	0.86		1.10
Pancreas	10.3	10.8	14.7	0.95		0.70
Cervix	9.7	7.0	9.5	1.39		1.02
Leukemia	9.1	11.1	8.8	0.82		1.03

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age. Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for the U.S. combined.

2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

3. Selected cancer sites by gender among the U.S. total population and the U.S. Hispanic/Latino population.

4. Abbreviations: NH - Non-Hodgkin; IBD - Intrahepatic Bile Duct NOS - Not Otherwise Specified

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIAv2) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates. Data Item 190 used in place of NHIA for MA and RI, see technical notes.