

Background

- Androgen deprivation therapy (ADT), i.e., hormonal therapy, has been the standard of care for men diagnosed with metastatic prostate cancer (mPC) since 1940.¹
- More recently, novel hormonal therapies and a combination therapy with ADT and a chemotherapy (e.g., docetaxel) with survival benefits have been available as well.^{1,2}
- A couple of studies using real-world data have reported underutilization of the novel treatments. But no population-based studies have examined treatment patterns in recent years to examine the potential impact of change in treatment landscape.^{3,4}
- Thus, we used data from a population-based registry to examine treatment utilization patterns in mPC.

Purpose

- To assess initial treatment patterns among men diagnosed with mPC.

Methods

- We used California Cancer Registry (CCR) data to identify men of ages ≥20 year, who were diagnosed with stage IV prostate cancer (site code: C619) between 2010–2018.
- We further restricted our sample to only first primary cases with microscopic confirmation and those with Gleason score ≥6.
- Treatment patterns were examined by assessing receipt of ADT and chemotherapy for initial cancer treatment.
- Frequency/proportion of patients who received ADT and/or chemotherapy were calculated by sociodemographic, comorbidity, and tumor characteristics.
- We, then, fitted a multinomial logistic regression model to examine whether receipt of ADT only and chemotherapy with or without ADT varied by sociodemographic and tumor characteristics after adjusting for covariates.


Results

- Of the 14,205 eligible men identified, most were ≥65 years old (64.2%) and non-Hispanic white (NHW, 58%), had Gleason score >7 (59%), and distant metastasis (68%, Table 1).
- In terms of treatments, the majority (n=9214, 64.9%) received ADT alone as initial cancer treatment, 1580 (11.1%) received both ADT and chemotherapy, 219 (1.5%) received only chemotherapy, 2942 (20.7%) received neither treatment (Table 1), and 250 (1.8%) had unknown treatment status (data not shown in Table 1).
- Our findings from regression model showed that patients of ages 65-74 and 85+ had significantly lower odds of receiving ADT (OR[95%CI]: 0.81[0.67, 0.98] and 0.62[0.49, 0.79] respectively) compared to 20-54 age group. On the other hand, all patients older than 54 were significantly less likely to receive chemotherapy +/- ADT (OR range: 0.04 in 85+ age group to 0.52 in 55-64 age group, p <0.0001 for all) compared to 20-54 age group (Figure 1).

Table 1. Type of initial treatment received by baseline sociodemographic and tumor characteristics.					
Characteristics	Total n (col %)*	Treatment Type, n (Row %)			
		Neither	Hormone Therapy (ADT)	ADT + Chemotherapy	Chemotherapy
Overall sample	14205 (100%)	2942 (20.71%)	9214 (64.86%)	1580 (11.12%)	219 (1.54%)
Age					
20-54	1068 (7.5%)	202 (18.91%)	615 (57.58%)	212 (19.85%)	22 (2.06%)
55-64	4019 (28.3%)	858 (21.35%)	2466 (61.36%)	551 (13.71%)	65 (1.62%)
65-74	4972 (35.0%)	1060 (21.32%)	3156 (63.48%)	566 (11.38%)	97 (1.95%)
75-84	2996 (21.1%)	540 (18.02%)	2164 (72.23%)	228 (7.61%)	27 (0.9%)
85+	1150 (8.1%)	282 (24.52%)	813 (70.7%)	23 (2%)	8 (0.7%)
Race/Ethnicity					
Non-Hispanic White	8283 (58.31%)	1721 (20.78%)	5334 (64.4%)	955 (11.53%)	124 (1.5%)
Non-Hispanic Black	1595 (11.23%)	348 (21.82%)	1047 (65.64%)	160 (10.03%)	20 (1.25%)
Hispanic	2888 (20.33%)	608 (21.05%)	1857 (64.3%)	307 (10.63%)	59 (2.04%)
Asian/Pacific Islander	1248 (8.79%)	223 (17.87%)	854 (68.43%)	135 (10.82%)	15 (1.2%)
Other/Unknown	191 (1.34%)	42 (21.99%)	122 (63.87%)	23 (12.04%)	1 (0.52%)
Neighborhood SES					
Lowest	2243 (15.79%)	500 (22.29%)	1431 (63.8%)	225 (10.03%)	46 (2.05%)
Lower-Middle	2597 (18.28%)	560 (21.56%)	1688 (65%)	262 (10.09%)	41 (1.58%)
Middle	3047 (21.45%)	657 (21.56%)	1963 (64.42%)	336 (11.03%)	37 (1.21%)
Upper-Middle	3131 (22.04%)	617 (19.71%)	2057 (65.7%)	368 (11.75%)	41 (1.31%)
Highest	2834 (19.95%)	542 (19.12%)	1839 (64.89%)	352 (12.42%)	49 (1.73%)
Clinical Gleason Score					
Gleason Score 6-7	2681 (18.87%)	887 (33.08%)	1551 (57.85%)	163 (6.08%)	19 (0.71%)
Gleason Score 8-10	8380 (58.99%)	1312 (15.66%)	5822 (69.47%)	1031 (12.3%)	90 (1.07%)
No needle biopsy/TURP performed	2256 (15.88%)	452 (20.04%)	1413 (62.63%)	301 (13.34%)	48 (2.13%)
Unknown	888 (6.25%)	291 (32.77%)	428 (48.2%)	85 (9.57%)	62 (6.98%)
Presence of any distant metastasis at diagnosis					
No distant metastasis	9671 (68.08%)	1600 (35.49%)	2610 (57.9%)	160 (3.55%)	37 (0.82%)
Distant metastasis	26 (0.18%)	1336 (13.81%)	6588 (68.12%)	1418 (14.66%)	180 (1.86%)
Unknown	4508 (31.74%)	6 (23.08%)	16 (61.54%)	2 (7.69%)	2 (7.69%)
Year of Diagnosis					
2010	1149 (8.09%)	260 (22.63%)	803 (69.89%)	59 (5.13%)	12 (1.04%)
2011	1176 (8.28%)	281 (23.89%)	814 (69.22%)	53 (4.51%)	14 (1.19%)
2012	1311 (9.23%)	320 (24.41%)	884 (67.43%)	56 (4.27%)	28 (2.14%)
2013	1385 (9.75%)	333 (24.04%)	952 (68.74%)	58 (4.19%)	15 (1.08%)
2014	1509 (10.62%)	313 (20.74%)	956 (63.35%)	168 (11.13%)	29 (1.92%)
2015	1677 (11.81%)	332 (19.8%)	1023 (61%)	262 (15.62%)	29 (1.73%)
2016	1865 (13.13%)	370 (19.84%)	1081 (57.96%)	345 (18.5%)	36 (1.93%)
2017	2006 (14.12%)	385 (19.19%)	1271 (63.36%)	285 (14.21%)	30 (1.5%)
2018	2127 (14.97%)	348 (16.36%)	1430 (67.23%)	294 (13.82%)	26 (1.22%)

*The total n and % do not add up to 100% due to exclusion of those with missing/unknown status.

Figure 1. Forest plots of adjusted odds ratios (ORs) and 95% confidence intervals for receipt of a) ADT only and b) Chemotherapy with or without ADT.



*The model was adjusted for following variables in addition to the ones listed above: Insurance type at diagnosis, baseline Charlson comorbidity score, and type of reporting facility (NCI, NCCN, CoC).

Results (contd.)

- Those residing in lower-to-middle SES neighborhoods had approximately 12% to 19% lower odds of receiving ADT. Although only findings for Lowest and Lower-to-middle SES groups were statistically significant (p-values: 0.009 and 0.03 respectively). While middle-to-lowest SES neighborhoods all had significantly lower odds of receiving chemotherapy +/- ADT (OR range: 0.60 - 0.70, p < 0.001) (Figure 1a and b).
- There was no difference in the receipt of ADT only by race/ethnicity, however, NHB men were significantly less likely to receive a chemotherapy +/- ADT compared to their NHW counterparts (0.76[0.61, 0.94]).
- Lastly, patients with Gleason score >7 and those with distant metastasis were significantly more likely to receive both ADT only and chemotherapy +/- ADT as initial cancer treatment.

Limitations

- Registry data do not include molecule level treatment information. Thus, we were unable to assess the use of novel ADTs and specific chemotherapies used.
- We were also unable to examine treatment received beyond the first course of treatment due to lack of data.

Conclusions

- Although ADT is the standard of care for mPC, about one-fifth of the patients still do not receive it.
- Use of combination therapy has increased since 2014. However, only a small portion of patients receive this treatment currently.
- There is some evidence of disparity in treatment utilization by age, neighborhood SES, and in NHB men for newly approved combination therapy.
- Further study is needed to confirm these disparities and to better understand reasons for existing treatment utilization.

Acknowledgments

This work was supported by the National Cancer Institute Surveillance, Epidemiology, and End Results Program (HHSN2612018000091).

References

1. Damodaran S, Kyriakopoulos CE, Jarrard DF. Newly Diagnosed Metastatic Prostate Cancer: Has the Paradigm Changed?. Urol Clin North Am. 2017;44(4):611-621. doi:10.1016/j.ucl.2017.07.008.
2. Mohler JL, Armstrong AJ, Bahnson RR, et al. Prostate Cancer, Version 1.2016. J Natl Compr Canc Netw. 2016;14(1):19-30. doi: 10.6004/jnccn.2016.0004.
3. Freedland SJ, Sandin R, Sah J, et al. Treatment patterns and survival in metastatic castration-sensitive prostate cancer in the US Veterans Health Administration. Cancer Med. 2021;10(23):8570-8580. doi: 10.1002/cam4.4372. Epub 2021 Nov 2.
4. Swami U, Sinnott JA, Haaland B, et al. Treatment Pattern and Outcomes with Systemic Therapy in Men with Metastatic Prostate Cancer in the Real-World Patients in the United States. Cancers. 2021; 13(19):4951. <https://doi.org/10.3390/cancers13194951>.